



Geometrical optics approximation for coefficients of expansion of phase function in Legendre polynomials

A.A. Kokhanovsky

Institute of Physics, 70 Skorina Ave.,
 Minsk 220072, Belarus

KEYWORDS

Radiative transfer theory, Phase function, Aerosols

The expanding of phase functions $x(\theta)$ of aerosol media in Legendre polynomials $P_l(\theta)$

$$x(\theta) = \sum_{l=0}^N x_l P_l(\theta) \quad (1)$$

is frequently used to solve the radiative transfer equation and to find radiative characteristics of aerosol layers.

The number of terms in eq. (1) is about 2ρ , where $\rho = ka$, $k = 2\pi/\lambda$, a is the radius of particle, λ is the wavelength. the coefficients x_l can be obtained from the following formula:

$$x_l = \frac{2l+1}{2} \int_0^\pi x(\theta) P_l(\theta) \sin \theta d\theta \quad (2)$$

There are some difficulties to use Eq. (2) for large ($a \gg \lambda$) particles, when one should calculate function $x(\theta)$ with the Mie theory. This problem can be simplified by using geometrical optics approximation for $x(\theta)$. The task of this paper is to derive simple formulae for calculation of the values of x_l within the framework of the geometrical optics approximation.

After substitution the geometrical optics approximation for the function $x(\theta)$ in Eq. (2) the following formula can be obtained:

$$x_l = \frac{\sigma_{sca}^d x_l^d + \sigma_{sca}^G x_l^G}{\sigma_{sca}^d + \sigma_{sca}^G} \quad (3)$$

where

$$\sigma_{sca}^d = \pi a^2, \quad \sigma_{sca}^G = \omega \pi a^2 \quad (4)$$

$$x_l^d = (2l+1)F(t), \quad x_l^G = (2l+1) \sum_{s=0}^l B_s \varphi_s(\alpha) \quad (5)$$

Here

$$\omega = \frac{1}{2} \sum_{j=1}^2 \int_0^{\pi/2} \left(R_j + \frac{(1 - R_j)^2 e^{-C\xi}}{1 - R_j e^{-C\xi}} \right) \sin 2\tau d\tau \quad (6)$$

$$F(t) = \frac{2}{\pi} (\arccos t - t\sqrt{1-t^2}) U_+(t), \quad B_s = \frac{\Gamma(s + \frac{1}{2}) \Gamma(l - s + \frac{1}{2})}{\pi \Gamma(s+1) \Gamma(l-s+1)} \quad (7)$$

$$\varphi_s(\alpha) = \int_0^{\pi/2} \frac{\Phi_{1s}(\alpha, \tau) + \Phi_{2s}(\alpha, \tau)}{2\omega} \sin 2\tau d\tau \quad (8)$$

$$\Phi_{js}(\alpha, \tau) = \frac{D_j(\alpha, \tau) + 2R_j^2 e^{-C\xi} (e^{-C\xi} - \cos 2\alpha\tau') \cos 2\alpha\tau}{1 - 2R_j e^{-C\xi} \cos(2\alpha\tau') + R_j^2 e^{-2C\xi}} \quad (9)$$

$$D_j(\alpha, \tau) = R_j(1 - e^{-2C\xi}) \cos 2\alpha\tau + (1 - R_j)^2 e^{-C\xi} \cos 2\alpha(\tau - \tau') \quad (10)$$

Γ is the gamma function, $U_+(t) = 0$ at $t > 1$ and $U_+(t) = 1$ at $t \leq 1$, $t = l/2\rho$, $c = 4\gamma\rho$, $\cos \tau' = \cos \tau/n$, $m = n - i\gamma$ is the complex refractive index of particles, $\xi = \sqrt{1 - \cos^2 \tau/n^2}$, R_j is the Fresnel reflection coefficients, $\alpha = l - 2s$.

Eq. (3) can be used to calculate the coefficients x_l for large ($\alpha \gg \lambda$) single particles. It was generalized on spherical polydispersions also. Note, that coefficients x_l^G decreases rapidly at $l \rightarrow \infty$ and $x_l^G \ll x_l^d$ as $l > 10$ for aerosol media. So at large l we have (see Eq. (3)) $x_l = x_l^d/(1 + \omega)$. The maximum of the function $x(l)$ is located at $l_m = \pi\rho/4$ and $x(l_m) = l_m/(1 + \omega)$.